

The change in oceanic O₂ inventory associated with recent global warming

Ralph F. Keeling[†] and Hernan E. Garcia[‡]

Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA 92093-0244

Communicated by Devendra Lal, University of California at San Diego, La Jolla, CA, March 15, 2002 (received for review September 24, 2001)

Oceans general circulation models predict that global warming may cause a decrease in the oceanic O₂ inventory and an associated O₂ outgassing. An independent argument is presented here in support of this prediction based on observational evidence of the ocean's biogeochemical response to natural warming. On time scales from seasonal to centennial, natural O₂ flux/heat flux ratios are shown to occur in a range of 2 to 10 nmol of O₂ per joule of warming, with larger ratios typically occurring at higher latitudes and over longer time scales. The ratios are several times larger than would be expected solely from the effect of heating on the O₂ solubility, indicating that most of the O₂ exchange is biologically mediated through links between heating and stratification. The change in oceanic O₂ inventory through the 1990s is estimated to be $0.3 \pm 0.4 \times 10^{14}$ mol of O₂ per year based on scaling the observed anomalous long-term ocean warming by natural O₂ flux/heating ratios and allowing for uncertainty due to decadal variability. Implications are discussed for carbon budgets based on observed changes in atmospheric O₂/N₂ ratio and based on observed changes in ocean dissolved inorganic carbon.

Repeated hydrographic surveys indicate that the upper 3 km of the oceans have warmed (1) and intermediate waters of high-latitude origin have freshened (2) over the past few decades. Model studies indicate that upper-ocean warming, high-latitude freshening, and an associated increase in the density stratification of the upper ocean are expected consequences of the changes in atmospheric radiative forcing caused by fossil-fuel burning and other human activities (3–5).

Repeated hydrographic surveys also indicate that small detectable changes have occurred in oceanic dissolved O₂ concentrations. As summarized in Table 1, detectable decreases in O₂ have been found in intermediate waters in the North Pacific, North Atlantic, South Pacific, and South Indian oceans, while small increases have possibly been found in deeper waters in the North Pacific and South Indian Oceans. What caused these O₂ changes is unclear, and different mechanisms, including changes in ocean circulation rates (6–9), changes in preformed values (10), changing Redfield ratios (11), and changes in biological production (8) have been offered as possible explanations in different regions. While the changes may partly reflect natural decadal variability, the clearest O₂ changes, found at intermediate depths, are in the direction of decreasing O₂ concentrations. A global reduction in dissolved O₂ is predicted by ocean general circulation models (OGCMs) driven by increasing greenhouse gases (12–15). In the model simulations, most of the O₂ decrease is attributed to enhanced stratification.

Stratification has two competing effects on subsurface oxygen concentrations. First, it reduces the upwelling of nutrients from deeper waters into surface waters, thus decreasing photosynthetic production and the associated flux of organic detritus into the ocean interior. This flux is often referred to as the “biological pump” (16), and reducing the rate of this pump increases subsurface O₂ concentrations by reducing subsurface O₂ utilization rates. Second, stratification limits the downward transport from O₂ from well-oxygenated surface waters into the ocean interior, which serves to reduce subsurface O₂ concentrations.

In the modeling studies cited above, the effect of stratification on O₂ transport exceeds the effect on subsurface O₂ utilization, leading to a net O₂ decrease. This result is expected, considering that stratification allows for more complete biological utilization of nutrients in surface waters, thus lowering the “preformed” (i.e., initial) nitrate and phosphate content of waters sinking into the oceans' interior. To conserve the total ocean nutrient inventory, an increase must occur in the inventory of nonpreformed nutrients—i.e., those that accumulate in subsurface waters from oxidative decomposition of organic detritus. Since O₂ is consumed by organic decomposition in proportion to the amount of nitrate or phosphate released (17), subsurface O₂ inventories can be expected to decrease in response to increased stratification. In general, the competing effects of the biological pump and vertical mixing on O₂ concentrations can be assessed based on their net impact on the vertical nutrient distributions. This net impact we refer to as the “efficiency” (as opposed to “rate”) of the biological pump.

Stratification can also be expected to induce a net release of O₂ from the ocean to the atmosphere. The oceanic O₂ inventory (I , mol) and sea-to-air O₂ flux (Z , mol yr⁻¹) are linked according to

$$dI/dt = -Z + \alpha dC_{\text{org}}/dt, \quad [1]$$

where C_{org} is the oceanic inventory of organic carbon, including both particulate and dissolved forms, and α is the O₂:C oxidative ratio for destruction or production of marine organic matter, and where small terms related to river and sediment transports have been neglected. The term $\alpha dC_{\text{org}}/dt$ effectively accounts for the column-integrated net production of O₂ by marine photosynthesis and respiration. This term is presumably small, although not necessarily negligible (18), because the main effect of marine photosynthesis and respiration on time scales of years to centuries is to redistribute inorganic materials within the ocean, rather than to cause accumulation or destruction of organic carbon (see also ref. 14). On these time scales, we therefore expect that changes in the oceanic O₂ inventory, due to stratification or other causes, should be roughly balanced by O₂ exchanges with the atmosphere ($dI/dt \approx -Z$).

Estimates based on OGCMs for oceanic O₂ outgassing due to global warming lie in the range of 0.2 to 0.7×10^{14} mol yr⁻¹ for the past few decades, with predictions of 1.0 to 1.6×10^{14} mol yr⁻¹ for late 21st century (12–15). According to these estimates, roughly a quarter of the predicted outgassing is attributable to the direct effect of warming on the O₂ solubility, while the remainder is due to increased stratification. The predicted changes for the past few decades amount to an average decrease in oceanic O₂ concentrations over a 20-year time frame of between 1 and 5 $\mu\text{mol kg}^{-1}$ if the changes are confined to the top 1,000 m of the oceans. Larger changes are predicted for the Southern Oceans due to reductions in deep convection (12, 13).

Abbreviations: OGCMs, ocean general circulation models; DIC, dissolved inorganic carbon.

[†]To whom reprint requests should be addressed. E-mail: rkeeling@ucsd.edu.

[‡]Present address: National Oceanographic and Atmospheric Administration, OCL-NODC, 1315 East-West Highway, Silver Spring, MD 20910.

Table 1. Significant recent changes in dissolved O₂ concentrations

Study	Location	Time span	Depth range, m	O ₂ change, μmol kg ⁻¹
Garcia <i>et al.</i> (10)	Atlantic 24°N section, averaged over full section	1981–1992	800–1,900	–3 to –7
Pahlow and Riebesell (11)	North Pacific, basin average	1950s–1990s	Unreported	~–5
Keller <i>et al.</i> (9)*	North Pacific, basin average	1970s–1990s	300–500	–1 to –2
			1,000–1,600	+2 to +3
Shaffer <i>et al.</i> (7)	Pacific, 28°S section, averaged from 52°–88°W	1967–1995	800–1,200	–5 to –8
Bindoff and McDougal (6)	Indian Ocean, 32°S section, averaged from 30°–117°E	1962–1997	300–800	–7 to –8
			2,500–4,000	+3
Matear <i>et al.</i> (13)	Southern Ocean, 110°–170°E, 50°–60°S	1965–1995	>400	–5 to –15
Emerson <i>et al.</i> (8)*	North Pacific, 154°W section, averaged from 22°–44°N	1980–1997	100–600	–9 to –20

Report change in O₂ – O₂^{}, rather than O₂ concentration, where O₂^{*} is the solubility.

It is hard to test these predictions with existing hydrographic data, given the sparse coverage and the lack of comprehensive syntheses.

There are several reasons why changes in the oceanic O₂ inventory could be important. Dissolved O₂ concentration is a useful diagnostic of ocean circulation and biological activity which can provide constraints on models of physical and biogeochemical response to climate change (13). Small changes in O₂ content could influence extent of hypoxic regions in coastal seas, in sediments, or in the open ocean, with consequences for the cycling of nitrogen and other redox-sensitive elements and for the distribution of many marine organisms (19). Changes in O₂ are diagnostic of changes in the efficiency of the marine biological pump, which may influence the rate at which the oceans absorb anthropogenic CO₂ (12). Finally, at least two approaches for estimating sinks of anthropogenic carbon dioxide require corrections for changes in oceanic O₂ inventory. The purpose of this paper is to provide an independent estimate of the plausible O₂ inventory changes associated with recent global warming and to discuss the implications for global carbon budgeting.

Natural Warming and O₂ Outgassing

It is easily established that a close association exists between ocean warming and O₂ outgassing over a range of space and time scales in the open ocean. At middle and high latitudes during the spring and summer, when the upper ocean is heated by the atmosphere, the oceans are a source of O₂ to the atmosphere, while in the fall and winter, when the upper ocean is cooled, the oceans are a sink of O₂ from the atmosphere. These seasonal air–sea O₂ exchanges are driven partly by biological exchanges, linked to seasonal stratification, nutrient supply, and irradiance, and partly by effects of heating and cooling on O₂ solubility (20).

We have estimated the ratio of seasonal O₂ outgassing to seasonal heating from global archived measurements of dissolved O₂ in surface waters, climatological winds, and climatological air–sea heat fluxes (21). The O₂ flux/heating ratio varies between 1.5 nmol of O₂ per joule at lower latitudes to 4 or 5 nmol of O₂ per joule in the 40° to 60° latitude bands, as shown in Fig. 1. Consistent with Najjar and Keeling (20), we find ratios that are larger than expected from the effect of warming on the O₂ solubility by factors between 1.5 and 2.5. We further have shown, based on comparisons with atmospheric O₂/N₂ data (21), that the component of the O₂ flux that correlates with heating dominates large-scale seasonal O₂ exchange. These results indicate that seasonal heating, through its effect on stratification, biological productivity, and O₂ solubility, is a major driver of the exchange, and not just coincidentally correlated with the exchange.

Linkages between heat fluxes and oxygen fluxes are also evident on longer time scales, as revealed from a plot of the tracer O₂^{*} = O₂ + 175PO₄ versus potential temperature (Θ), as

shown in Fig. 2. The tracer O₂^{*}, which is identical to Broecker's tracer PO₄^{*} (25) but expressed in O₂ rather than PO₄ units, is a measure of the O₂ gained or lost by a water parcel through air–sea gas exchange (26). O₂^{*} is largely conserved below the sea surface, where photosynthesis and respiration produce compensating effects on O₂ and PO₄. O₂^{*} keeps track of air–sea O₂ exchanges driven by both solubility changes and the processes controlling the efficiency of the biological pump. Assuming rapid air–sea equilibration, the solubility component is simply given by the O₂ solubility (O₂^{*}), so that the remaining biological pump component is proportional to the (apparent) preformed phosphate content (PO₄ – (O₂^{*} – O₂)/175) of the water (26).

A prominent feature in Fig. 2 is the strong association between O₂^{*} and Θ in waters of the main thermocline between 6° and 18°C. Waters around 18°C consistently have lower O₂^{*} than waters around 6°C, indicating that conversion of cooler water to warmer water, which occurs mostly at low latitudes, leads to outgassing of O₂, while the conversion of warmer water back to cooler water, which occurs mostly at higher latitudes, leads to uptake of O₂. Outside the North Atlantic, the O₂^{*}/Θ slope of ~22 μmol kg⁻¹ °C⁻¹ in waters between 6° and 18°C, is equivalent to ~5 nmol of O₂ per joule of warming, as derived




Fig. 1. Ratio of the seasonal O₂ outgassing to the seasonal upper ocean warming by 10° latitude bands. Dashed line indicates the ratios expected based on changes in O₂ solubility alone. Ratios from 20°N to 20°S are poorly defined due to weak seasonal forcing. The seasonal O₂ outgassing here accounts for the anomalous flux after the mean annual outgassing or ingassing is subtracted.




Fig. 2. Scatter plot of the tracer $O_2^* = O_2 + 175PO_4$ versus potential temperature for data collected on the Geochemical Ocean Section Study (22–24). The data through the main thermocline of the Indian and Pacific oceans are seen to scatter around a line with a slope of approximately $-22 \mu\text{mol kg}^{-1} \text{ } ^\circ\text{C}^{-1}$. Changes in O_2^* are generally larger than expected from changes in the O_2 solubility (dashed line). Antarctic surface water data are from ref. 25. NADW, North Atlantic deep water.

by multiplying by seawater density and dividing by heat capacity. The slope is several times larger than expected from solubility changes alone, indicating that the O_2 exchanges are mainly controlled by variations in the efficiency of the marine biological pump. A generally weaker and but more variable O_2 /heat relationship is indicated for waters warmer than about 18°C , which is consistent with the lower nutrient content of these waters and a reduction of influence of biological relative to solubility effects.

A less steep O_2^*/θ trend is seen in Fig. 2 for the thermocline of the North Atlantic Ocean compared with other oceans, possibly owing to the lower nutrient content of the North Atlantic. The North Atlantic trend of $\sim 13 \mu\text{mol kg}^{-1} \text{ } ^\circ\text{C}^{-1}$ (equivalent to $\sim 3 \text{ nmol J}^{-1}$) connects low-latitude surface waters in the Atlantic with North Atlantic Deep Water (NADW), which lies below the O_2^*/θ trend of the other oceans. A steeper O_2^*/θ ratio of around 30 to $40 \mu\text{mol kg}^{-1} \text{ } ^\circ\text{C}^{-1}$ (equivalent to $7.5\text{--}10 \text{ nmol J}^{-1}$) is found for the deep Antarctic sequence (25), which is driven by ventilation of deep waters around Antarctica. Here stratification induced by warming and freshening in the summer months inhibits the uptake of O_2 by deeper waters, while the breakdown of stratification induced by wintertime cooling and brine rejection from sea ice enhances O_2 uptake and deepwater formation (27). The air–sea exchanges and water-mass-mixing around Antarctica effectively convert circumpolar deep water, which derives largely from NADW, into colder Antarctic surface waters. By this conversion, the oceans around Antarctica release heat to the atmosphere and take up O_2 .

Due to sparse coverage, Fig. 2 does not resolve well water masses in the Equatorial Pacific. Here, however a very different relationship between heating and O_2 flux is known to exist. Equatorial upwelling raises cool, oxygen-deficient waters to the surface, where a net uptake of O_2 from the atmosphere (20) and a net heating of the water occurs. The O_2 flux/heating ratio in the Equatorial Pacific is thus opposite in sign to the cases considered above. This feature is a result of the upwelled waters having been exposed to the surface for a very brief period of weeks or days, which is insufficient to allow O_2 to equilibrate with the atmosphere and for biological production to remove the nutrients. As the upwelled waters spread laterally away from the Equator, the net effect of warming and nutrient withdrawal leads to an overall O_2 release to the atmosphere (20, 28). Integrated over a wider latitude band, the net effect of Equatorial upwelling on heat and O_2 exchange is therefore more concordant with the main thermocline trend.

The patterns noted above suggest the following generalizations: Although the changing efficiency of the biological pump dominates the O_2 response of the ocean to warming and stratification, the O_2 response is nevertheless strongly tied, over a range of space and time scales, to the net air–sea heat flux. For time scales of months to centuries, the O_2 flux/heating ratios generally lie in the range of 2 to 10 nmol J^{-1} . Larger ratios are found at higher latitudes, particularly in the Southern Hemisphere and for processes occurring over time scales of decades to centuries (e.g., thermocline ventilation) compared with time

Table 2. Estimated oceanic O₂ outgassing 1990–2000 due to changing radiative forcing

Ocean region	Heating		O ₂ /heat ratio, nmol J ⁻¹	O ₂ outgassing	
	10 ²² J yr ⁻¹	%		10 ¹⁴ mol yr ⁻¹	%
North Atlantic	0.18 ± 0.08	30	3 ± 1.5	0.05 ± 0.04	18
Deep Southern Ocean	0.06 ± 0.03	10	10 ± 5.0	0.06 ± 0.04	20
Other oceans	0.36 ± 0.16	60	5 ± 2.5	0.18 ± 0.12	61
Global total	0.6 ± 0.18	100	4.9	0.29 ± 0.13	100

Heating rates are based on observed warming in each region from 1955 to 1995 (1), scaled to yield total of 0.6×10^{22} J yr⁻¹, the warming attributable to increasing radiative forcing from 1990 to 2000 (4, 5). Error analysis assumes 45% uncertainty in individual heating rates and 50% uncertainty in O₂/heat ratios, with all errors uncorrelated.

scales of months (seasonal exchanges). Some differences exist from ocean to ocean, with the North Atlantic having smaller O₂ response per unit heating or cooling than other high-latitude regions. This analysis does not resolve the O₂ response of the oceans to heating and cooling on thousand-year and longer time scales.

Anthropogenic Warming and O₂ Outgassing

In Table 2 we use the O₂ flux/heating relationships found above to formulate an estimate of the global air–sea O₂ flux from 1990 to 2000. We consider three ocean regions: (i) North Atlantic (at all depths), (ii) the deep Southern Ocean (>1,000 m), and (iii) the remaining oceans (at all depths). Warming in each region is assumed to produce O₂ outgassing proportional to observed steady-state O₂/Θ relationships in these regions. The approach is motivated by the rough universality of O₂/heat ratios for processes ranging from warming and cooling on seasonal time scales to steady-state warming/cooling over decades to centuries. The approach effectively adopts the null hypothesis that the ocean’s response to transient warming on decadal time scales is governed by similar ratios.

Because compilations of ocean warming (1) have been completed only through year 1998, and because these compilations do not resolve warming on a yearly basis below 300-m depth, we rely on model simulations (4, 5) as a means of extrapolating the observed long-term (1950s–1990s) warming in each region through the 1990–2000 period. The approach effectively accounts for the warming caused by anthropogenic radiative forcing but not natural variability. The latter we treat as a source of noise. We derive a central estimate of the global O₂ outgassing of 0.29×10^{14} mol of O₂ yr⁻¹ for 1990–2000, where 18% of the total O₂ outgassing is attributable to warming in the North Atlantic, 20% in the deep Southern Ocean, and 61% in the remainder of the upper ocean. Our estimate corresponds to a global average O₂ flux heating ratio of 5 nmol J⁻¹.

The uncertainty around our central estimate attributable to uncertainties in regional O₂/Θ ratios and long-term warming rates is $\pm 0.13 \times 10^{14}$ mol of O₂ yr⁻¹. A much larger uncertainty must be allowed for decadal variability (15, 29, 30). One estimate of this can be derived by multiplying the global O₂/heat ratio of 5 nmol of O₂ J⁻¹ by the decadal variability in global ocean heat storage, which we estimate from heat storage data (1), after removing the long-term trend, to be $\pm 5 \times 10^{22}$ J, which yields $\pm 2.5 \times 10^{14}$ mol of O₂ variability on a decadal basis. Here we adopt a slightly higher estimate of $\pm 4 \times 10^{14}$ mol of O₂, on the grounds that the heat storage data (1) may underestimate true variability due to spatial and temporal averaging, and given uncertainties in the appropriate O₂/heat ratio. Treating the decadal variability as a source of random noise, we derive an estimate of $0.29 \pm 0.4 \times 10^{14}$ mol of O₂ yr⁻¹ for the total oceanic O₂ outgassing from 1990 to 2000.

From 1990 to 1998, the global upper ocean (<300 m) heat content increased at a rate of $\sim 0.5 \times 10^{22}$ J yr⁻¹ (1), which is faster than the rate of $\sim 0.2 \times 10^{22}$ J yr⁻¹ that we would estimate

from projecting long-term warming rates. If we assume that the additional warming of $\sim 0.3 \times 10^{22}$ J yr⁻¹ persisted through year 2000, and we add the difference to our outgassing estimate, then our central estimate increases from 0.29×10^{14} mol yr⁻¹ to 0.44×10^{14} mol yr⁻¹, assuming a scaling of 5 nmol J⁻¹. To apply such a correction is premature, however, because the global 1990–1998 heating trend is heavily influenced by a large anomaly in the North Atlantic in 1998, which was possibly a transient associated with the 1997–1998 El Niño event (1). In any case, the correction would be within our allowed uncertainties for decadal variability of $\pm 0.4 \times 10^{14}$ mol yr⁻¹. Once heat storage data are compiled through 2000, it may be possible to refine the decadal outgassing estimate and reduce the allowed uncertainties.

Carbon Budgeting

What are the implications of oceanic O₂ outgassing for carbon budgeting? A correction for oceanic O₂ outgassing is needed to estimate land and ocean carbon sinks based on the global budgets of atmospheric O₂ and CO₂ (31, 32). These can be written

$$\Delta\text{CO}_2 = F - O - B \quad [2]$$

and

$$\Delta\text{O}_2 = -\alpha_F F + \alpha_B B + Z, \quad [3]$$

where F is the CO₂ source from fossil-fuel burning, O and B are, respectively, the ocean and land sinks, α_F and α_B are global-average O₂:C exchange ratios for fossil-fuel and land biota, and Z is the net source of O₂ from the oceans. Eqs. 2 and 3 are solved to yield estimates of O and B , using ΔO_2 and ΔCO_2 derived, respectively, from observed changes in atmospheric O₂/N₂ ratio and CO₂ mole fraction, F and α_F derived from industrial records, and $\alpha_B \approx 1.1$ (31, 32).

Previously it has been assumed that the ocean outgassing term Z is zero to within the uncertainties (31, 32), or allowance has alternately been made for O₂ outgassing based on the solubility effect alone (33). In Table 3, we correct the estimate of Manning (33), as cited in the recent Intergovernmental Panel on Climate Change (34) report, based on our above estimate of $Z = 0.29 \pm 0.4$ mol of O₂ per yr, where this estimate implicitly allows for the effect of anthropogenic warming on both solubility and stratification. The correction increases the oceanic sink by 0.18 Pg of C yr⁻¹ and decreases the land sink by the same amount relative to the Manning (33) estimate. Although the change is small relative to other uncertainties, it nevertheless helps to reconcile the estimated oceanic sink with recent model estimates (35).

A correction for oceanic O₂ is also needed for carbon budgets based on direct measurement of changes in dissolved inorganic carbon (DIC) in the ocean. Detecting the input of anthropogenic CO₂ into the oceans is difficult due to large natural variability in DIC caused by ocean biology. This difficulty is commonly overcome by normalizing to a constant O₂ concentration to filter out the variability due to ocean

Table 3. O₂-based global carbon budget 1990–2000

Budget item	Pg of C yr ⁻¹	
	Manning (33)	Revised budget
Fossil-fuel burning	6.33 ± 0.4	6.33 ± 0.4
Atmospheric CO ₂ increase	3.21 ± 0.1	3.21 ± 0.1
Total ocean CO ₂ sink	1.68 ± 0.5	1.86 ± 0.6
Total land CO ₂ sink	1.44 ± 0.7	1.26 ± 0.8

Budgets are based on the decadal trend in atmospheric O₂/N₂ ratio at two northern hemisphere stations in the Scripps O₂/N₂ network. Manning (33) assumes $Z = 0.16 \pm 0.16 \times 10^{14}$ mol of O₂ yr⁻¹; revised budget assumes $0.29 \pm 0.4 \times 10^{14}$ mol of O₂ yr⁻¹ (see text). Budgets differ in assumed ocean warming rate, and in assumed oceanic O₂ outgassing per unit warming. Both budgets allow for outgassing of N₂ (which affects the observed O₂/N₂ ratio) at a rate based on N₂ solubility of 2.2 nmol of N₂ per J of warming.

biology (36–40). In effect, what is reported is not the change DIC, but rather the change in the quantity DIC + O₂/α, where α ≈ 1.3 (36, 37). By design, this approach neglects changes in DIC caused by variations in the efficiency of the biological pump. If the dissolved O₂ inventory decreases globally due to stratification, the approach will underestimate oceanic CO₂ uptake by an amount given approximately by (dI/dt)/α, where dI/dt is the change in oceanic O₂ inventory. The correction is the same sign and a similar magnitude to that required based on atmospheric O₂ and CO₂ budgets. Taking the observed ocean warming of ~2 × 10²³ J between the middle 1950s and 1990s (1), and assuming the O₂ inventory decreases by 5 nmol of O₂ J⁻¹, yields an upwards correction of ~9 Pg of C (1 Pg = 10¹⁵ g) over the 1955–1995 period. In comparison, the total oceanic uptake from preindustrial times through 1990 is estimated to be 107 ± 27 Pg of C yr⁻¹ (34).

Recent estimates of change in oceanic inventories of DIC (38–40) have not allowed for changing O₂ inventory, and therefore are presumably biased low, although more work is needed to establish reliable corrections in the individual ocean basins and globally. If the DIC data are additionally normalized based on regressions against Θ and alkalinity, then additional corrections may be needed for changing ocean heat content (independent of the effect on O₂) and alkalinity inventory.

Discussion

Our estimate of 0.29 × 10¹⁴ mol of O₂ yr⁻¹ for recent oceanic O₂ outgassing caused by changing radiative forcing lies within the range estimated based on recent OGCMs (12–15). The relatively good agreement between these independent estimates strengthens the case that a long-term ocean outgassing of this magnitude may actually be occurring. Nevertheless, the similarity in these estimates undoubtedly results partly from similar assumptions. For example, the estimates commonly assume that Redfield P/O₂ ratios are constant with time and space, and that phosphate is the ultimate limiting nutrient. Also, while our estimate assumes that the O₂ response to transient warming is similar to the O₂ response to steady-state warming and cooling, the OGCMs similarly assume that the response of the ocean biology to transient warming can be understood on the basis of parameters adjusted to duplicate steady-state behavior. It is unclear if these assumptions form a realistic basis for projection.

Our estimate effectively assumes that O₂/Θ relationships—or equivalently, preformed phosphate/Θ relationships—remain constant during transient warming. For example, the O₂ flux/

heating ratio we adopt for warming of the main thermocline would be obtained if the main effect of warming was to deepen the thermocline without altering the preformed phosphate/Θ relationship across the thermocline, or if the surface isotherms progress polewards during the transient at the same rate as the isolines of surface phosphate. The O₂ flux/heating ratio we adopt for the deep Southern Ocean would be obtained if warming caused a reduction in deepwater formation around Antarctica, thus increasing the influence of other source waters on the chemistry of the deep ocean.

Our estimate of the effect of Southern Ocean warming on O₂ exchange can be compared with the study Broecker *et al.* (41), who used PO₄³⁻, ¹⁴C, and chlorofluorocarbon (CFC) data to support the argument that the ventilation rate of deep waters around Antarctica slowed substantially in the 20th century. According to their scenario, the PO₄³⁻ content of the deep ocean should now be decreasing with time, in which case a net O₂ outgassing should also be occurring, as required by the links between PO₄³⁻ (i.e., O₂^{*}) and the air–sea O₂ flux. Taking their estimate of the difference in PO₄³⁻ content of southern source waters relative to average deep waters and their estimate that the input rate has slowed by 10 × 10⁶ m³ s⁻¹ yields a required outgassing rate of 0.3 × 10¹⁴ mol of O₂ yr⁻¹, which is 5 times larger than our estimate of 0.06 × 10¹⁴ mol of O₂ yr⁻¹ for the deep Southern Ocean (Table 2). We defend our smaller estimate on the following grounds: First, the Broecker *et al.* scenario implies that the deepwater heat content should be increasing by ~0.4 × 10²² J yr⁻¹, based on the ~3°C difference in temperature between southern surface waters and average deep waters. A warming rate of this magnitude is inconsistent with the observed temperature trends (1), unless the warming is mostly confined below 3,000 m, the maximum depth considered by Levitus *et al.* (1). Second, recent work (42) suggests that a large 20th-century slow-down is not necessary to explain the CFC and PO₄³⁻ data.

A reliable assessment of the global air–sea O₂ flux will ultimately require an approach based on direct observations rather than model studies. Over the next few decades, global systematic decreases in dissolved O₂ can be expected at the level of 0.4 × 10¹⁴ mol of O₂ yr⁻¹ or larger, which corresponds to a change of 0.7 μmol of O₂ kg⁻¹ per decade, if spread uniformly over 2,000 m. Resolving these changes against natural variability will require a high measurement density, but this may be feasible with appropriate sensor and platform development and with a concerted long-term observing program. Resolving these changes is needed for carbon budgeting, as a complement to lower-density DIC measurements and atmospheric O₂/N₂ measurements, and would help to assess the overall impact of climate change on the biogeochemistry and biodiversity of the oceans. Another source of uncertainty in carbon budgets involves change in oceanic organic carbon (18), which we have assumed is small, but is not well constrained. We suggest that a program to directly monitor oceanic inventories of O₂ and organic carbon, along with inorganic carbon and nutrients, should be given some priority in future ocean observing systems.

We thank Scott Doney, Ray Najjar, and Corinne Le Quéré for helpful comments. This work was supported by the National Science Foundation under Grant ATM-0000923, National Oceanic and Atmospheric Administration under Grant NA77RJ0453A, and the National Aeronautics and Space Administration under Grant NAG5–6668, and was completed in part while one of us (R.K.) was hosted at the Max Planck Institute for Biogeochemistry in Jena, Germany.

- Levitus, S., Antonov, J. I., Boyer, T. P. & Stephens, C. (2000) *Science* **287**, 2225–2229.
- Wong, A. P. S., Bindoff, N. L. & Church, J. A. (1999) *Nature (London)* **400**, 440–443.

- Manabe, S., Stouffer, R. J., Spelman, M. J. & Bryan, K. (1991) *J. Climate* **4**, 785–818.
- Levitus, S., Antonov, J. I., Wang, J., Delworth, T. L., Dixon, K. W. & Broccoli, A. J. (2001) *Science* **292**, 267–270.

5. Barnett, T. P., Pierce, D. W. & Schnur, R. (2001) *Science* **292**, 270–274.
6. Bindoff, N. L. & McDougall, T. J. (2000) *J. Phys. Oceanogr.* **30**, 1207–1222.
7. Shaffer, G., Leth, O., Ulloa, O., Bendtsen, J., Daneri, G., Dellarossa, V., Hormazabal, S. & Sehlstedt, P.-I. (2000) *Geophys. Res. Lett.* **27**, 1247–1250.
8. Emerson, S., Mecking, S. & Abell, J. (2001) *Global Biogeochem. Cycles* **15**, 535–554.
9. Keller, K., Slater, R. D., Bender, M. & Key, R. M. (2002) *Deep-Sea Res. II* **49**, 345–362.
10. Garcia, H., Cruzado, A., Gordon, L. & Escanez, J. (1998) *J. Geophys. Res.* **103**, 2817–2830.
11. Pahlow, M. & Riebesell, U. (2000) *Science* **287**, 831–833.
12. Sarmiento, J. L., Hughes, T. M. C., Stouffer, R. J. & Manabe, S. (1998) *Nature (London)* **393**, 245–249.
13. Matar, R. J., Hirst, A. C. & McNeil, B. I. (2000) *Geochem. Geophys. Geosyst.* **1**, American Geophysical Union paper no. 10.1029/2000GC000086.
14. Plattner, G.-K., Joos, F., Stocker, T. F. & Marchal, O. (2001) *Tellus Ser. B* **53**, 564–592.
15. Bopp, L., Le Quéré, C., Heimann, M., Manning, A. C. & Monfray, P. (2002) *Biogeochem. Cycles*, in press.
16. Volk, T. & Hoffert, M. I. (1985) in *The Carbon Cycle and Atmospheric CO₂ Archean to Present*, AGU Geophysical Monograph 32, eds. Sundquist, E. T. & Broecker, W. S. (Am. Geophys. Union, Washington, DC), pp. 99–110.
17. Redfield, A. C., Ketchum, B. H. & Richards, F. A. (1962) in *The Sea*, ed. Hill, M. N. (Interscience, New York), Vol. 2, pp. 26–77.
18. Hansell, D. A. & Carlson, C. A. (2001) *Deep-Sea Res. II* **48**, 1649–1667.
19. Rogers, A. D. (2000) *Deep-Sea Res. II* **47**, 119–148.
20. Najjar, R. G. & Keeling, R. F. (2000) *Global Biogeochem. Cycles* **14**, 573–584.
21. Garcia, H. E. & Keeling, R. F. (2001) *J. Geophys. Res.* **106**, 31155–31160.
22. Bainbridge, A. E. (1981) *Hydrographic Data 1972–1973, GEOSECS Atlantic Expedition* (National Science Foundation, Washington, DC), Vol. 1.
23. Broecker, W. S., Spenser, D. W. & Craig, H. (1982) *Hydrographic Data 1973–1974, GEOSECS Pacific Expedition* (National Science Foundation, Washington, DC), Vol. 3.
24. Weiss, R. F., Broecker, W. S., Craig, H. & Spenser, D. W. (1983) *Hydrographic Data 1977–1978, GEOSECS Indian Ocean Expedition* (National Science Foundation, Washington, DC), Vol. 5.
25. Broecker, W. S., Peacock, S. L., Walker, S., Weiss, R., Fahrback, E., Schroeder, M., Mikolajewicz, U., Heinze, C., Key, R., Peng, T.-H. & Rubin, S. (1998) *J. Geophys. Res.* **103**, 15833–15843.
26. Keeling, R. F. & Peng, T.-H. (1995) *Philos. Trans. R. Soc. London B* **348**, 133–142.
27. Gordon, A. L. & Huber, B. A. (1990) *J. Geophys. Res.* **95**, 11655–11672.
28. Stephens, B. B., Keeling, R. F., Heimann, M., Six, K. D., Murnane, R. & Caldeira, K. (1998) *Global Biogeochem. Cycles* **12**, 213–230.
29. Bender, M., Ellis, T., Tans, P., Francey, R. & Lowe, D. (1996) *Global Biogeochem. Cycles* **10**, 9–21.
30. McKinley, G. A., Follows, M. J. & Marshall, J. (2000) *Geophys. Res. Lett.* **27**, 2933–2936.
31. Keeling, R. F. & Shertz, S. R. (1992) *Nature (London)* **358**, 723–727.
32. Battle, M., Bender, M. L., Tans, P. P., White, J. W. C., Ellis, J. T., Conway, T. & Francey, R. J. (2000) *Science* **287**, 2467–2470.
33. Manning, A. C. (2001) Ph.D. thesis (Univ. of California, San Diego).
34. Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Khashgi, H. S., Le Quéré, C., Scholes, R. J., Wallace, D. W. R., et al. (2001) in *Climate Change 2001, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Pitelka, L. & Ramirez Rojas, A. (Cambridge Univ. Press, Cambridge, U.K.), pp. 183–237.
35. Orr, J. C., Maier-Reimer, E., Mikolajewicz, U., Monfray, P., Sarmiento, J. L., Toggweiler, J. R., Taylor, N. K., Palmer, J., Gruber, N., Sabine, C. L., et al. (2001) *Global Biogeochem. Cycles* **15**, 43–60.
36. Brewer, P. G. (1978) *Geophys. Res. Lett.* **5**, 997–1000.
37. Chen, C.-T. A. & Millero, F. J. (1979) *Nature (London)* **277**, 205–206.
38. Gruber, N. (1998) *Global Biogeochem. Cycles* **12**, 165–191.
39. Sabine, C. L., Key, R. M., Johnson, K. M., Millero, F. J., Poisson, A., Sarmiento, J. L., Wallace, D. W. R. & Winn, C. D. (1999) *Global Biogeochem. Cycles* **13**, 179–198.
40. Peng, T.-H., Wanninkhof, R., Bullister, J. L., Feely, R. A. & Takahashi, T. (1998) *Nature (London)* **396**, 560–563.
41. Broecker, W. S., Sutherland, S. & Peng, T.-H. (1999) *Science* **286**, 1132–1135.
42. Orsi, A. H., Jacobs, S. S., Gordon, A. L. & Visbeck, M. (2001) *Geophys. Res. Lett.* **28**, 2923–2926.